





令和8年度 フェロモントラップ調査結果(シロイチモジヨトウ)

※調査データはグラフの下にあります。





調査データ

地帯区分		中東部			中東部			北西部			南西部		
設置場所		三次市三和町			世羅郡世羅町山中福田			安芸高田市高宮町			東広島市安芸津町		
周辺作物		アスパラガス			ねぎ			ねぎ			ばれいしょ		
月	半旬	本年	平均(2年)	前年	本年	平均(2年)	前年	本年	平均(5年)	前年	本年	平均(1年)	前年
5月	1	0	0.2	0	0	1.5	3.0	0	4.1	0.8	0	1.3	1.3
	2	0	0.3	0	0	1.5	3.0	0	2.6	2.3	0	1.0	1.0
	3	0	1.4	0	0.3	1.5	3.0	0	2.5	2.3	0.6	0.7	0.7
	4	0	2.3	1.4	1.4	1.5	3.0	0	7.9	0	3.1	2.1	2.1
	5	0	2.8	0.3	1.4	0.6	1.2	2.3	15.3	3.3	2.5	2.1	2.1
	6	0.6	2.2	0	0.9	1.3	1.7	2.7	11.2	0.7	6.3	3.1	3.1
6月	1	1.4	2.6	0	0	3.0	5.9	8.1	2.9	0	11.5	3.6	3.6
	2	1.4	3.1	4.7	0	5.8	11.5	8.4	4.2	0	11.5	3.6	3.6
	3	1.9	4.8	16.3	1.7	6.2	12.4	8.8	6.2	0	5.3	6.6	6.6
	4	3.3	4.2	7.0	3.9	6.6	11.5	15.8	6.1	0.6	8.6	8.0	8.0
	5	7.9	7.8	22.7	6.4	5.7	8.9	17.5	11.0	1.4	28.0	5.7	5.7
	6	2.7	6.1	3.6	12.1	2.9	4.3	3.5	10.7	0	25.2	5.1	5.1
7月	1	3.1	5.9	1.9	5.7	4.9	5.4	0	11.4	0	13.8	4.4	4.4
	2	5.7	5.6	1.9	1.7	4.6	4.9	0	14.1	0	9.6	3.6	3.6
	3	3.6	6.4	2.5	2.9	3.6	3.6	0	20.4	0	6.8	6.4	6.4
	4		5.6	1.2	1.7	2.6	3.8		23.7	0.1		3.0	3.0
	5		6.9	2.0		2.9	4.8		28.3	0.7		0.9	0.9
	6		7.3	2.3		6.0	7.5		16.7	0.9		1.7	1.7
8月	1		7.4	3.0		11.7	13.3		13.5	0.7		0.7	0.7
	2		10.8	25.6		10.0	15.8		16.1	10.9		20.0	20.0
	3		10.5	22.0		7.2	11.9		18.8	24.4		28.0	28.0
	4		10.6	15.0		6.1	4.9		20.3	6.3		8.6	8.6
	5		10.4	8.6		17.5	4.4		18.0	6.3		5.1	5.1
	6		22.1	31.0		21.5	9.1		36.4	10.6		7.3	7.3
9月	1		25.0	19.0		17.2	10.0		32.5	13.4		20.1	20.1
	2		13.1	23.4		15.8	10.6		29.0	21.4		69.3	69.3
	3		15.3	43.6		14.3	12.1		27.8	28.3		90.4	90.4
	4		21.4	53.7		20.4	16.4		27.8	30.0		138.9	138.9
	5		19.4	62.2		20.6	22.1		33.6	14.0		203.6	203.6
	6		22.3	85.8		20.7	33.3		45.3	10.0		561.4	561.4
10月	1		35.1	207.9		26.1	44.8		32.9	10.0		558.4	558.4
	2		27.9	156.4		23.7	41.2		20.4	2.6		460.0	460.0
	3		16.0	99.1		7.5	10.0		21.7	0.7		74.3	74.3
	4		3.4	7.1		6.8	4.3		22.3	0.7		25.1	25.1
	5		1.6	1.4		3.8	1.7		22.0	0.7		14.3	14.3
	6		0.6	0		1.4	0		23.0	0.9		22.4	22.4